



Children's Feel Better Cold and Fever Suspension



Ibuprofen ► Pain reliever/
(NSAID) Fever reducer

Pseudoephedrine HCL ►
Nasal decongestant

Relieves:
Fever,
Sore Throat
Stuffy Nose

Grape Flavor



4 Fl Oz (120 mL)

Drug Facts

Active ingredients

(in each 5 ml teaspoonful)
Ibuprofen 100 mg (NSAID)*.....Pain reliever/
fever reducer
Pseudoephedrine HCl 15mg.....Nasal decongestant
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves
- nasal and sinus congestion
- stuffy nose
- head ache
- minor body aches and pains
- sore throat
- fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

Sore throat warning: Severe or persistent sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer and/or nasal decongestant
- right before or after heart surgery
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI), certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- problems or serious side effects from taking pain relievers, fever reducers, or nasal decongestants
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- taken a diuretic

Ask a doctor or pharmacist before use if the child is

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any continuing medical condition
- taking any other drug

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- the child does not get any relief during the first day (24 hours) of treatment
- fever, pain, or nasal congestion gets worse or lasts for more than 3 days
- stomach pain or upset gets worse or lasts
- symptoms continue or get worse
- redness or swelling is present in the painful area
- the child gets nervous, dizzy, or sleepless
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- shake well before using
- find right dose on chart below. If possible use weight to dose; otherwise use age.
- use only with enclosed measuring cup
- if needed, repeat dose every 6 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yrs)	Dose (teaspoonful)
under 24	under 2	ask a doctor
24-47	2 to 5	1 tsp
48-95	6 to 11	2 tsp

Other information

- do not use if printed neckband is broken or missing
- store at 15-30°C (59-86°F)
- see bottom of box for lot number and expiration date

Inactive ingredients acesulfame potassium, citric acid, corn syrup, D&C red #33, FDA&C blue #1, FD&C red #40, flavors, glycerin, hypromellose, polysorbate 80, purified water, sodium benzoate, xanthan gum

Questions? 1-800-123-4567

<http://ChildrensFeelBetter.com>